



# Exploring the diversity of microorganisms and potential pectinase activity isolated from wet fermentation of coffee in northeastern Peru

Samia Littly Jahavelly FERNANDEZ-GÜIMAC<sup>1</sup> , Jhordy PEREZ<sup>1</sup> , Jani Elisabet MENDOZA<sup>1</sup> ,  
Danilo Edson BUSTAMANTE<sup>1,2</sup> , Martha Steffany CALDERON<sup>1,2\*</sup> 

## Abstract

In this study, the microbiota associated with coffee fermentation from two regions from northern Peru was evaluated. Bacteria and fungi were isolated from coffee farms in the Amazonas and Cajamarca regions and identified using molecular markers 16S rRNA and ITS. The potential pectinase activity was registered by the formation of a transparent halo around colonies. As a result, 29 and 09 species belonging to bacteria and fungi, respectively, were found. The bacterial genera *Lysinibacillus* and *Stenotrophomonas* and the fungal genus *Aspergillus* accounted for the highest number of species isolated from coffee ferments. Forty-one out of 71 isolates showed some type of pectinase enzyme activity, and they included 23 isolates from Cajamarca and 18 from Amazonas. Nevertheless, only three bacterial species registered the formation of transparent halos and showed relevant potential pectinase enzyme activity, namely, *Lysinibacillus xylanilyticus*, *Stenotrophomonas maltophilia*, and *Stenotrophomonas pavanii*, which were all from the Cajamarca region. These species could be further investigated by quantifying enzymes activity and performing other biochemical properties to prototype starter cultures. Accordingly, the study of indigenous microorganisms with biological potential will be essential to increase the coffee value chain and improve the incomes of farmers.

**Keywords:** coffee; indigenous microorganisms; northern Peru; potential pectinase activity; wet coffee processing.

**Practical Application:** *Lysinibacillus xylanilyticus*, *Stenotrophomonas maltophilia*, and *S. pavanii* showed potential pectinase activity.

## 1 Introduction

Coffee is one of the most popular nonalcoholic beverages (Haile & Kang, 2019) and has been considered the most traded and consumed food product in the world for decades (Silva et al., 2013). The coffee fruit consists of an outer layer of skin (exocarp) that covers a layer of pulp followed by mucilage (mesocarp), which is firmly attached to the rigid layer called parchment (endocarp) (Evangalista et al., 2015). Coffee fruits undergo fermentation processes under either wet, dry or semidry conditions (Evangalista et al., 2014; Schwan et al., 2012). During fermentation, the mucilaginous layer is removed and special characteristics are conferred regarding its flavor and aroma because of the production of a wide range of metabolites, such as organic acids, alcohols and esters, which will later add complexity to coffee (De Bruyn et al., 2016; Feng et al., 2016; Vera et al., 2018).

Although the main characteristic flavor of coffee comes from the chemical composition of the bean, some of the metabolites that contribute to the sensory characteristics of the beverage are produced by microbiota associated with the fermentation process (Silva et al., 2013). This microbiota is mainly composed of filamentous fungi, yeasts, archaea, and lactic and acetic acid bacteria (Silva et al., 2005; Huch & Franz, 2015). The main techniques characterizing these microorganisms include culture-

independent and culture-dependent approaches (Duong et al., 2020). Culture-independent approaches determine the relative abundance of microorganisms (metabarcoding), potential function of the associated genes (metagenomics), and identification of microorganisms (barcodes), mainly based on DNA techniques (Feng et al., 2016; Oliveira et al., 2013; Santos et al., 2020). Among the main DNA markers used in these approaches are 16S, 18S or 26S/28S rDNA for bacteria (Martins et al., 2020) and ITS or *tef1* for fungi (Bustamante et al., 2019). The amplification and sequencing of these markers have greatly increased the number of identified microbial species (Duong et al., 2020).

Conversely, the culture-dependent strategy allows the characterization of microorganisms individually on the basis of their morphology and biochemical and functional traits, including their pectinolytic activity (Abdollahzadeh et al., 2020; de Melo-Pereira et al., 2014; Teshome et al., 2017). Pectinase enzymes catalyze the degradation of pectins of the coffee mucilaginous layer by depolymerization and deesterification reactions (Wood & Kellogg, 1988; Xavier-Santos et al., 2004). Pectinase is a general term referring to the mixture of three different enzymatic activities, namely, polygalacturonase (PG), pectin esterase (PE) and pectin lyase (PL), which contribute to the breakdown and modification of pectins and the release of compounds, such as

Received 10 Sept., 2022

Accepted 14 Nov., 2022

<sup>1</sup>Instituto de Investigación para el Desarrollo Sustentable de Ceja de Selva – INDES-CES, Universidad Nacional Toribio Rodríguez de Mendoza, Amazonas, Peru

<sup>2</sup>Facultad de Ingeniería Civil y Ambiental, Universidad Nacional Toribio Rodríguez de Mendoza, Amazonas, Peru

\*Corresponding author: martha.calderon@untrm.edu.pe

galacturonic acid and its oligomers, during the fermentation process (Combo et al., 2012; Patidar et al., 2018; Sunnotel & Nigam, 2002). Currently, the application of pectinolytic enzymes plays an important role in food technology, especially in the fermentation process (Poveda et al., 2018; Souza et al., 2003). These enzymes provide opportunities for future research on the development of a starter culture from microorganisms with pectinolytic activity (Silva et al., 2013). It is important to note that culture-independent and culture-dependent approaches have their own strengths and weaknesses; however, both are of the utmost importance for microorganism analyses, and better results are achieved if they work in a complementary manner (Duong et al., 2020).

Because coffee producers in northern Peru do not employ standardized techniques, the duration of the fermentation process varies among coffee farms (10–48 hours), resulting in a wide range of coffee grades (Puerta & Echeverry, 2015). This scenario might involve a diversity of microorganisms with specific pectinolytic activity during the fermentation process of coffee in northern Peru. Therefore, the aim of this study was to characterize the diversity of microorganisms associated with spontaneous coffee wet fermentation from two regions (Amazonas and Cajamarca) in northeastern Peru through the culture of microorganisms. For this purpose, bacteria and fungi were isolated from coffee farms and identified using Sanger sequencing of molecular markers 16S rRNA for bacteria and ITS for fungi. Additionally, potential pectinase activity was evaluated to list the species that could be subsequently studied and develop a starter inoculum in fermentation to improve coffee quality.

## 2 Materials and methods

### 2.1 Sample collection

A total of twenty-four samples of coffee beans under fermentation at different times (from 5 to 32 hours of fermentation) were collected from coffee farmers in the Amazonas and Cajamarca regions. Accordingly, a Drigalski spatula was dipped into the fermented beans and sown directly into Petri dishes with 39 g/L potato dextrose agar medium (PDA, HIMEDIA) (Silva et al., 2013). Petri dishes were sealed with parafilm, labeled and then transported to the Laboratory of Molecular Biology and Genomics of the Universidad Nacional Toribio Rodríguez de Mendoza under sterilized conditions and stored at 28 °C for seven days under 12 h photoperiods (white fluorescent light/darkness) (da Silva et al., 2021).

### 2.2 Bacterial and fungal isolation

After colony formation, microorganisms were isolated on selective media according to Garofalo et al. (2015). Briefly, ~0.1 g of grown colonies was transferred to De Man, Rogosa and Sharpe (MRS, HIMEDIA) agar media containing 1200 µl/L fluconazole (Elhalis et al., 2020; Oktaviani et al., 2020) for bacterial isolation and yeast extract peptone dextrose (YPD) agar media containing 600 µl/L azithromycin for fungal isolation (de Melo-Pereira et al., 2014; Evangelista et al., 2014). MRS and YPD Petri dishes were incubated at 30 °C with 12 h photoperiods (white fluorescent

light/dark) for bacterial (for 48–72 hours) and fungal growth (for 5–7 days) (Elhalis et al., 2020; Roussos et al., 1995).

### 2.3 DNA sequencing

Isolates were superficially scraped from Petri dishes with a sterilized scalpel and placed in pre-labeled 2.0 mL Safelock Eppendorf tubes. Genomic DNA was extracted using a DNA Miniprep Kit (ZymoBIOMICS, California, USA) following the manufacturer's instructions. Two markers were sequenced for bacterial (16S ribosomal RNA, 16S rRNA) and fungal (internal transcribed spacer, ITS) identification. The primer combination for the 16S rDNA gene was 16S\_341F: 5'-CCTACGGGAGGCAGCAG-3'; 16S\_800R: 5'-CAGGACTACCAGGGTATCTAAT-3' (Lechner et al., 1998) and for ITS1-5.8S, ITS1\_F: 5'-TCCGTAGGTGAACCTGCGG-3'; ITS2\_R: 5'-TCTCCTCCGCTTATTTGATATATGC-3' (Masoud et al., 2004). Each marker was amplified using polymerase chain reaction (PCR) with MasterMix (Promega, Madison, Wisconsin, USA) in the following reaction mixture: 10 ng of DNA and 0.25–0.5 pmol of the forward and reverse primers in a total volume of 10 µL. The PCR protocol for 16S rRNA followed a pre-denaturation step (95 °C for 3 min); 35 cycles of denaturation (94 °C for 1 min), annealing (55 °C for 1 min), and extension (72 °C for 1 min); and a final elongation step (72 °C for 5 min). Additionally, the PCR protocol for ITS followed a pre-denaturation step (94 °C for 7 min); 40 cycles of denaturation (94 °C for 45 sec), annealing (57 °C for 45 sec), and extension (72 °C for 1 min); and a final elongation step (72 °C for 7 min). Amplicons were purified using a Macherey-Nagel Kit (NucleoSpin Gel and PCR Clean-up, Düren, Germany). Sequences of the forward and reverse strands were commercially determined by Macrogen (Seoul, South Korea) and then edited in Chromas v1.45 software (Technelysium, 1998). A total of 71 sequences were generated and deposited in GenBank (Table S1).

### 2.4 Phylogenetic analyses

Sequences were initially aligned using MUSCLE algorithms (Thompson et al., 1994) and manually adjusted with MEGA7 (Kumar et al., 2016). In total, eight phylogenetic trees corresponding to the class level were generated for bacteria (Betaproteobacteria, Gammaproteobacteria, Alphaproteobacteria, Firmicutes, and Actinobacteria) and fungal groups (Eurotiomycetes, Sordariomycetes, and Dothideomycetes) on the basis of 16S rRNA and ITS markers, respectively. The nucleotide substitution model that best matched each marker and group was selected using PartitionFinder (Lanfear et al., 2012) (Table S2). The best partitioning strategy and the best sequence evolution model were selected based on the Bayesian information criterion (BIC). Maximum likelihood (ML) analyses were performed with raxmlHPC-AVX (Stamatakis, 2014) implemented in the raxmlGUI 1.3.1 interface (Silvestro & Michalak, 2012) using the best model as appropriate with 1000 bootstrap replicates for both datasets. Bayesian inference (BI) was carried out with MrBayes 3.2.6 (Ronquist et al., 2012) using Metropolis coupled Markov chain Monte Carlo (MCMC) and the best model as appropriate for both datasets. We conducted two runs with four chains each (three hot chains and one cold chain) for 10000000 generations, and trees were sampled every 1000 generations. We plotted the

probability versus generation using Tracer 1.6.6 (Bouckaert et al., 2014) until a probability plateau was reached and set the burn-in value (Bustamante et al., 2021). Intraspecific and interspecific pairwise divergence was estimated using the p-distance model in MEGA7.

### 2.5 Potential pectinase activity

Potential pectinolytic activity was determined using the methods described by Hankin & Lacy (1984) and Schwan et al. (1997). Isolated yeasts and bacteria were cultured at 30 °C for four days with 12 h photoperiods on plates with mineral medium containing 5 g/L polygalacturonic acid (MP5) at pH 5.5 for detecting polygalacturonase (PG) activity and mineral medium containing 5 g/L pectin (MP7) at pH 7.2 for detecting pectin lyase (PL) activity (Hankin & Lacy, 1984). The mineral medium contained 5 g/L glucose, 6 g/L  $\text{KH}_2\text{PO}_4$ , 1 g/L yeast extract, 2 g/L  $(\text{NH}_4)_2\text{SO}_4$ , 15 g/L agar and 0.1 ml/L solutions of 0.1 g/L  $\text{FeSO}_4$ , 20 g/L  $\text{MgSO}_4$ , 0.1 g/L  $\text{CaCl}_2$ , 0.2 g/L  $\text{H}_3\text{BO}_3$ , 0.2 g/L  $\text{MnSO}_4$ , 1.4 g/L  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ , 1 g/L  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ , and 0.2 g/L  $\text{MoO}_3$  (Carrim et al., 2006; Silva et al., 2008a). Potential enzymatic activity was registered by the formation of a transparent halo around colonies due to the cell lysis and precipitation of polysaccharides during enzymatic reactions with pectin after the addition of 1% cetyl trimethylammonium bromide (CTAB) (Wang & Stegemann, 2010; Zhao et al., 2015). Briefly, 5 mL of CTAB solution was added to each plate and incubated at room temperature for three hours in dark conditions (García et al., 2007; García & Chafra, 2015). Afterward, the excess CTAB solution was removed. Six measurements for the area, perimeter, length, and diameter of halos were recorded from each culture using ImageJ software (Huang et al., 2007).

## 3 Results

### 3.1 Phylogenetic analyses

A total of 71 microorganisms were isolated from coffee fermentation processes in northeastern Peru. Fifty-eight (81.69%) isolates were bacterial, and thirteen (18.31%) isolates were fungal. Twenty-four bacterial and 5 fungal isolates were identified from Amazonas, whereas 34 bacterial and 8 fungal isolates were identified from Cajamarca (Figure S1). The genus *Lysinibacillus* was the most abundant bacteria (28%), and the genus *Aspergillus* (46%) was the dominant fungi. Additionally, 29 species of bacteria from Amazonas (15 species) and Cajamarca (19 species) were identified and distributed in 14 genera. The most frequent genera in Amazonas were *Lysinibacillus* (46%) and *Bacillus* (13%), whereas such genera in Cajamarca were *Stenotrophomonas* (29%) and *Lysinibacillus* (14%). Nine fungal species were identified and distributed in seven genera. The most frequent genera in Amazonas were *Aspergillus* (60%), *Fusarium* (20%), and *Metarhizium* (20%), whereas those in Cajamarca were *Aspergillus* (38%), *Epicoccum* (25%), and *Penicillium* (13%). Eight phylogenetic trees were constructed using 16S rRNA for bacteria (258 sequences) and ITS for fungi (125 sequences). Species identity was assigned based on grouping with sequences of type species and low values of genetic divergence. In either case, we followed a conservative approach where not every lineage was recognized as a species.

### 3.2 Bacterial groups

#### Class Actinobacteria

The phylogeny of the class Actinobacteria included species of the genus *Streptomyces* (29 sequences, 1451 bp) (Figure S2). *Kitasatospora kazusensis* Li was used as an outgroup. In this class, *Streptomyces* sp. (MET119) was recognized in samples from Cajamarca, and it was sisters to *S. coelestis* Krassilnikov and *S. violaceoruber* Waksman & Curtis. The low genetic divergences among these species confirm no variations between sequences of type sequences (i.e., *S. violaceoruber*- *S. coelestis*; *S. niveoruber*- *S. griseoviridis*).

#### Class Alphaproteobacteria

The phylogeny of the class Alphaproteobacteria included members of the genera *Pseudochrobactrum* and *Ochrobactrum* (29 sequences, 1429 bp) (Figure S3). *Alcaligenes faecalis* Castellani & Chalmers was used as the outgroup. Three species were recognized from Amazonas and three from Cajamarca. *P. asaccharolyticum* Kämpfer (MET110), *Pseudochrobactrum* sp.1 (MET10), and *Pseudochrobactrum* sp.2 (MET116) were identified. The last two species differed by 0.3% in genetic variation. Other *Pseudochrobactrum* species ranged from 0 to 4.2%. Additionally, *Ochrobactrum* showed a paraphyletic relationship and low divergence between its species. For instance, type specimens of *O. tritici* Leubner and *O. cystisi* Zurdo-Piñero varied by 0.1%. *Ochrobactrum pseudogrignonense* Kämpfer (MET88), *Ochrobactrum* sp.1 (MET71 and MET115), and *Ochrobactrum* sp.2 (MET19) were identified. The genetic divergence between *Ochrobactrum* sp. 1 and *Ochrobactrum* sp. 2 was 1.5%.

#### Class Betaproteobacteria

The phylogeny of the class Betaproteobacteria included members of the genera *Alcaligenes*, *Achromobacter*, and *Delftia* (42 sequences, 1491 bp) that were grouped into three different clades (Figure S4). *Paenibacillus amylolyticus* Nakamura was used as an outgroup. One species was recognized from Amazonas and five from Cajamarca. Clade A contained three specimens of *Alcaligenes* sp. with identical sequences (MET72, MET85, and MET106). The genetic divergence between the type sequences of *A. aquatilis* van Trappen and *A. faecalis* Castellani & Chalmers was 0.5%. In clade B, *Achromobacter spiritinus* Vandamme (MET81 and MET117) and *Achromobacter* sp. (MET20) were found and differed by 1.3%. Genetic divergences between type sequences of *Ach. animicus* Vandamme - *Ach. mucicolens* Vandamme and *Ach. ruhlandii* Packer and Vishniac - *Ach. denitrificans* Rügner & Tan were 0.4% and 0.5%, respectively. Genetic variation in *Ach. spiritinus* Gomila lineage ranged from 0.1-0.3%. In clade C, *Delftia lacustris* Jørgensen (MET12, MET80) and *Delftia* sp. (MET23) were found. Genetic divergences between our *Delftia* specimens were 2.4%, whereas the type sequences of *D. tsuruhatensis* Shigematsu - *D. litopenaei* Chen and *D. acidovorans* den Dooren de Jong - *D. litopenaei* Chen differed by 2.0% and 2-2.6%, respectively.

#### Class Firmicutes

The phylogeny of the class Firmicutes included members of the genera *Alkalihalobacillus*, *Bacillus*, *Clostridium*, *Lysinibacillus*, and *Paenibacillus* (109 sequences, 1529 bp) (Figure S5). *Proteus*



*hauseri* O'Hara was used as the outgroup. Six species were recognized from Amazonas and eight from Cajamarca. In clade A, *Clostridium algidixylanolyticum* Broda (MET11) was identified. This species was in sistership to *C. aerotolerans* van Gylswyk & van der Toorn. In clade B, *Paenibacillus campinasensis* Yoon (MET87) and *Paenibacillus* sp. (MET22) were identified. The latter was closely related to *P. mobiliz* Yang. In clade C, the following species were identified: *Bacillus cereus* Frankland & Frankland (MET112 and MET113), *B. graminis* Bibi (MET109), *B. subtilis* Ehrenberg (MET68), *Lysinibacillus fusiformis* Priest (MET78, MET83, MET86, MET92, MET93, MET97, MET98, MET107, MET111, MET120, and MET118), *L. xylanilyticus* Lee (MET9), *Lysinibacillus* sp.1 (MET8 and MET17), and *Lysinibacillus* sp.2 (MET84 and MET114). The genetic divergence between *Lysinibacillus* sp. 1 and *Lysinibacillus* sp. 2 ranged from 0.6-0.8%. In this clade, *A. clausii* Nielsen (MET13) and *Alkalihalobacillus* sp. (MET64 and MET6) were also identified, and they differed by 4.6-5.3%.

#### Class Gammaproteobacteria

The phylogeny of the class Gammaproteobacteria included members of the genera *Morganella*, *Proteus*, and *Stenotrophomonas* (49 sequences, 1436 bp) (Figure S6). *Clostridium algidixylanolyticum* Broda was used as an outgroup. Five species were recognized from Amazonas and two from Cajamarca. In clade A, *Morganella* sp. (MET91), *P. columbae* Dai (MET89), and *Proteus* sp. (MET90) were identified. *Morganella* sp. differed from *M. morganii* Winslow by 0.3-0.7%. In clade B, *Stenotrophomonas maltophilia* Hugh (MET5, MET65, MET66, MET69, and MET70) and *S. pavanii* Ramos (MET1, MET2, MET4, MET7, MET18, MET67, and MET77) were identified. The genetic variation between *S. maltophila* and *S. pavanii* was 1.2%.

### Fungi

#### Class Dothideomycetes

The phylogeny of the class Dothideomycetes included members of the genera *Cladosporium* and *Epicoccum* (37 sequences, 539 bp) (Figure S7). *Aspergillus creber* Jurjević, S.W. Peterson & B.W. Horn was used as the outgroup. Two species were recognized from Cajamarca. *Cladosporium* sp. (MET131) was identified in clade A, while *Epicoccum ovisporum* Valenz-Lopez, Stchigel, Crous, Guarro & J.F. Cano (MET105 and MET132) was placed in clade B.

#### Class Eurotiomycetes

The phylogeny of the class Eurotiomycetes included members of the genera *Aspergillus*, *Penicillium*, and *Talaromyces* (51 sequences, 732 bp) (Figure S8). *Fusarium keratoplasticum* Geiser, O'Donnell, D.P.G. Short & Ning Zhang was used as the outgroup. Two species were recognized from Amazonas and four from Cajamarca. Clade A contained *Talaromyces cnidii* T.S.H. Yu, T.J. An & H.K. Sang (MET104), while clade B contained *Penicillium citrinum* Thom (MET126). Clade C comprised *Aspergillus creber* Jurjević, S.W. Peterson & B.W. Horn (MET122 and MET123) and two unidentified species, *Aspergillus*

sp.1 (MET127 and MET128) and *Aspergillus* sp.2 (MET121 and MET129). *Aspergillus* sp.1 differed by 0.5% from *Aspergillus* sp.2.

#### Class Sordariomycetes

The phylogeny of the class Sordariomycetes included members of the genera *Fusarium* and *Metarhizium* (37 sequences, 564 bp) (Figure S9). *Beauveria bassiana* (Bals.-Criv.) Vuill. was used as the outgroup. Two species were recognized from Amazonas. Clade A contained *Fusarium* sp. (MET103), whereas clade B contained *Metarhizium guizhouense* Chen & Guo (MET130). Genetic variation within *M. guizhouense* ranged from 0.2-1.5%.

### 3.2 Potential pectinase activity

The 71 isolated microorganisms were cultured in the media types MP5 (with pectin) and MP7 (with polygalaturonic acid) to perform an initial screening of the pectinase enzyme activity. After adding 1% CTAB to the plates and incubating for three hours, the formation of halos around the growing colonies was evaluated. In total, 41 isolates showed the formation of different types of halos around the colony in MP5 (35 isolates) and MP7 (24 isolates) media (Table 1, Figure 1, Figure S10). Eighteen isolates from Amazonas and 23 from Cajamarca showed halos. However, only three species (i.e., *Lysinibacillus xylanilyticus*, *Stenotrophomonas maltophilia*, and *S. pavanii*) formed transparent halos of 4.16-7.55 cm (Table 2, Figure 2).

## 4 Discussion

In this study, the diversity of the microbial community in coffee wet fermentation was evaluated under culture approaches using Sanger sequencing of molecular markers for identification (i.e., 16S rRNA and ITS) while an initial screening of the pectinase activity was performed (i.e., MP5 and MP7 media). The higher number of identified species and isolates with PG and PL activities were found in Cajamarca region, which typically uses a longer fermentation time and produces higher coffee quality than the Amazonas region (Agencia Peruana de Noticias, 2019). Several bacteria and fungi identified in this study were previously reported in other coffee fermentation processes. For instance, bacterial species such as *B. cereus*, *B. subtilis*, *L. fusiformis*, and *Paenibacillus* sp. from Minas Gerais, Brazil (Evangelista et al., 2015; Silva et al., 2008a,b) and *Proteus* sp. from Mysore, India (Agate & Bhat, 1966) were isolated from either wet or semidry fermentation processes. On the fungal side, members of *Aspergillus*, *Fusarium*, and *Penicillium* were isolated from Brazil and Colombia (Cruz-O'Byrne et al., 2021; Silva et al., 2000) from wet coffee fermentation. The identification of these bacteria and fungi in our study suggests that they are commonly isolated in specific Petri dishes (Hamdouche et al., 2016; Vilela et al., 2010). Conversely, this is the first time that the following microorganisms were isolated from coffee ferment in either bacterial or fungal media, *Alkalihalobacillus*, *Pseudochrobactrum*, *Stenotrophomonas*, and *Talaromyces*, confirming the importance of this approach to characterize the diversity of microorganisms that has not been recorded previously in fermentation processes (Carvalho et al., 2017; Elhalis et al., 2020).

**Table 1.** Isolates from Amazonas (A) or Cajamarca (C) Regions that showed halo formation around the colony in MP5 and MP7 media.

Species Bacteria	Code	Region	Time*	Media	Features of halo
<i>Bacillus cereus</i>	MET112	A	5	MP5	White color, non-uniform intensity, no definite shape
<i>Lysinibacillus fusiformis</i>	MET78	A	5	MP5	Whitish color, uniform intensity, continuous rounded shape
<i>Lysinibacillus fusiformis</i>	MET97	A	5	MP5, MP7	Both: beige in color, uniform intensity, continuous rounded shape
<i>Lysinibacillus fusiformis</i>	MET98	A	5	MP5	White color, uniform intensity, discontinuous rounded shape
<i>Lysinibacillus fusiformis</i>	MET111	A	5	MP5	Yellowish white color, uniform intensity, continuous rounded shape
<i>Lysinibacillus</i> sp.2	MET84	A	5	MP5	Yellowish white color, uniform intensity, continuous rounded shape
<i>Pseudochrobactrum asaccharolyticum</i>	MET110	A	5	MP5	White color, uniform intensity, no definite shape
<i>Pseudochrobactrum</i> sp.2	MET116	A	5	MP7	Beige color, uniform intensity, continuous rounded shape
<i>Alcaligenes</i> sp.	MET106	C	6	MP7	Beige color, uniform intensity, continuous rounded shape
<i>Clostridium algidixylanolyticum</i>	MET11	C	6	MP5	Yellowish white color, non-uniform intensity, discontinuous rounded shape
<i>Delftia lacustris</i>	MET12	C	6	MP5, MP7	MP5: yellowish white color, MP7: beige color, both: uniform intensity, continuous rounded shape
<i>Delftia lacustris</i>	MET80	C	6	MP5	Yellowish white color, uniform intensity, continuous rounded shape
<i>Paenibacillus</i> sp.	MET22	C	6	MP5	Beige color, uniform intensity, continuous rounded shape
<i>Stenotrophomonas maltophilia</i>	MET69	A	9	MP5	Beige color, uniform intensity, continuous rounded shape
<i>Stenotrophomonas pavanii</i>	MET18	A	9	MP7	Yellowish white color, non-uniform intensity, discontinuous rounded shape
<i>Lysinibacillus fusiformis</i>	MET86	A	10	MP5, MP7	MP5: yellowish white color, MP7: beige color with yellowish border, both: non-uniform intensity, discontinuous rounded shape
<i>Lysinibacillus fusiformis</i>	MET83	C	12	MP5, MP7	In MP5: yellowish white color, in MP7: yellowish beige color, Both: uniform intensity, continuous rounded shape
<i>Ochrobactrum</i> sp.2	MET19	C	12	MP5	White color, non-uniform intensity, discontinuous rounded shape
<i>Achromobacter spiritinus</i>	MET81	C	18	MP5, MP7	MP5: Transparent white color, MP7: Beige color, both: uniform intensity, continuous rounded shape
<i>Delftia</i> sp.	MET23	C	18	MP5, MP7	Both: beige-yellowish color, uniform intensity, continuous rounded shape
<i>Lysinibacillus fusiformis</i>	MET107	C	22	MP5	Yellowish white color, uniform intensity, continuous rounded shape
<i>Stenotrophomonas maltophilia</i>	MET65	C	22	MP5, MP7	MP5: yellowish white color, in MP7: transparent color with beige gradient edges, Both: uniform intensity, continuous rounded shape.
<i>Stenotrophomonas pavanii</i>	MET1	C	22	MP5, MP7	MP5: yellowish-white color, MP7: transparent, brownish-white gradient edges, Both: uniform intensity, continuous rounded shape
<i>Stenotrophomonas pavanii</i>	MET2	C	22	MP5	Brownish beige color, uniform intensity, non-continuous rounded shape
<i>Bacillus graminis</i>	MET109	C	27	MP7	Beige color, uniform intensity, continuous rounded shape
<i>Lysinibacillus</i> sp.1	MET8	C	27	MP5	Yellowish white color, uniform intensity, continuous rounded shape
<i>Morganella</i> sp.	MET91	A	27	MP5, MP7	MP5: white color, MP7: beige color, Both: uniform intensity, continuous rounded shape
<i>Proteus</i> sp.	MET90	A	27	MP5, MP7	MP5: beige color, MP7: white color, Both: uniform intensity, continuous rounded shape.
<i>Stenotrophomonas maltophilia</i>	MET70	C	27	MP5, MP7	MP5: yellowish white color, MP7: pinkish color with beige gradient edges, Both: uniform intensity, continuous rounded shape.

\* Fermentation time (5h, 6h, 9h, 10h, 12h, 18h, 22h, 27h, 32h) at the moment of collection.

**Table 1.** Continuação...

Species Bacteria	Code	Region	Time*	Media	Features of halo
<i>Stenotrophomonas pavanii</i>	MET7	C	27	MP7	Yellowish-white color, non-uniform intensity, continuous rounded shape
<i>Stenotrophomonas pavanii</i>	MET77	C	27	MP5, MP7	MP5: yellowish white color, MP7: pinkish color with beige gradient edges, Both: uniform intensity, continuous rounded shape
<i>Alcaligenes</i> sp.	MET72	C	32	MP5, MP7	MP5: yellowish white color, MP7: beige color with yellowish edges, Both: uniform intensity, continuous rounded shape
<i>Lysinibacillus fusiformis</i>	MET92	A	32	MP5, MP7	In MP5: White color, in MP7: Beige color, both with non-uniform intensity and discontinuous rounded shape.
<i>Lysinibacillus fusiformis</i>	MET93	A	32	MP5	Yellowish white color, non-uniform intensity, continuous rounded shape
<i>Lysinibacillus xylanilyticus</i>	MET9	C	32	MP5, MP7	MP5: yellowish white color, MP7: transparent color with beige edges, Both: uniform intensity, continuous rounded shape
<i>Lysinibacillus</i> sp.1	MET17	A	32	MP5, MP7	MP5: beige color, non-uniform intensity, MP7: yellowish white color, uniform intensity, Both: continuous rounded shape
<i>Ochrobactrum</i> sp.1	MET71	C	32	MP5	MP5: yellowish color, MP7: yellowish white color, Both: uniform intensity, continuous rounded shape
<i>Pseudochrobactrum</i> sp.1	MET10	C	32	MP7	Yellowish-white color, non-uniform intensity, no definite shape
<b>Fungi</b>					
<i>Penicillium citrinum</i>	MET126	C	12	MP5, MP7	Both: beige in color, uniform intensity, continuous rounded shape
<i>Aspergillus</i> sp.1	MET127	C	22	MP5, MP7	MP5: beige color with yellowish edges, MP7: pink color with beige edges, Both: uniform intensity, continuous rounded shape
<i>Talaromyces cnidii</i>	MET104	C	27	MP5	Beige color, uniform intensity, continuous rounded shape

\* Fermentation time (5h, 6h, 9h, 10h, 12h, 18h, 22h, 27h, 32h) at the moment of collection.

**Table 2.** Measurements of halos and colonies of selected microorganisms with pectinolytic activity.

Code	Genus	Halo Diameter (HD)	Colony Diameter (CD)	HD/CD Ratio
MET65	<i>Stenotrophomonas maltophilia</i>	5.71	3.63	1.57
MET1	<i>Stenotrophomonas pavanii</i>	7.55	5.04	1.50
MET9	<i>Lysinibacillus xylanilyticus</i>	4.16	2.42	1.72


The most common microorganisms reported in coffee fermentation are lactic acid bacteria (LAB), such as *Bacillus*, *Lactobacillus*, *Lactococcus*, *Leuconostoc*, and *Weissella* (Melo-Pereira et al., 2017; Pothakos et al., 2020), whereas the most commonly reported fungal species belonged to *Candida*, *Hanseniaspora*, *Pichia*, and *Saccharomyces* (Oliveira-Junqueira et al., 2019; Zhang et al., 2019). However, the predominant species in this study corresponded to the bacteria *Lysinibacillus* and the fungi *Aspergillus*. This highlights the importance of region-specific evaluations that allow the characterization of indigenous microbiota in coffee fermentation processes (Carvalho et al., 2017; Elhalis et al., 2020). This might be a consequence of specific environmental factors (e.g., temperature and altitude) (Evangelista et al., 2015; Körner, 2007).

Furthermore, some of the identified species have been associated with good coffee quality, such as *Cladosporium cladosporioides* (Chalfoun et al., 2007; Pereira et al., 2005), while others were linked to low quality, such as *Fusarium* and *Penicillium* species


(Masoud & Kaltoft, 2006; Souza et al., 2017) and some *Aspergillus* species, which represent potential ochratoxin A producers (Joosten et al., 2001; Mantle & Chow, 2000). However, once fermentation is finished, their abundances are not significant, ensuring that the final coffee is safe for human consumption consume (Masoud & Kaltoft, 2006; Souza et al., 2017).

Out of 71 isolated microorganisms, 41 showed potential pectinolytic enzyme activity (pectin degradation) on the basis of their PG (85.4%) and PL (58.5%) and combined PG and PL (43.9%) activities. PG and PL enzymes have been extensively reported in species of the genera *Aspergillus*, *Bacillus* and *Penicillium* (Chaudhri & Suneetha, 2012; Kashyap et al., 2000; Murad & Azzaz, 2011). This study confirms potential PG and PL activities in these three and 12 other genera (Table 1). Pectinase enzyme activity is associated with the presence of transparent halos due to the formation of alcohols and acids (e.g., butyric acetic, lactic and other long-chain carboxylic acids) (Barragán et al., 2014; Beg et al., 2000; Carrim et al., 2006; García & Chafla, 2015).





**Figure 1.** Formation of halos in MP5 and/or MP7. Microorganisms isolated from the Cajamarca Region: *Penicillium citrinum* (A, MP5), *Delftia lacustris* (B, MP5), *Aspergillus* sp.1 (C, MP5; D, MP7), *Ochrobactrum pseudogrignonense* (E, MP5), *Stenotrophomonas pavanii* (F, MP5), *Alcaligenes* sp. (G, MP7), *Clostridium algidixylanolyticum* (H, MP5). Microorganisms isolated from Amazonas: *Lysinibacillus fusiformis* (I, MP7; J, MP7; K, MP5; L, MP5).



**Figure 2.** Microorganism colonies before and after precipitating 1% CTAB on MP7 media and forming transparent halos. A: *Stenotrophomonas maltophilia* without CETAB, A': *Stenotrophomonas maltophilia* with CETAB, B: *Stenotrophomonas pavanii* without CETAB, B': *Stenotrophomonas pavanii* with CETAB, C: *Lysinibacillus xylanilyticus* without CETAB, C': *Lysinibacillus xylanilyticus* with CETAB.

This scenario was reported exclusively in MP5 medium for the following three bacterial species *Lysinibacillus xylanilyticus* (MET9), *Stenotrophomonas maltophilia* (MET65), and *S. pavanii* (MET1). These species showed higher halo diameters ( $5.8 \pm 1.7$  cm) than those reported in other species ( $2.33 \pm 0.4$  cm and  $1.6 \pm 0.82$  cm), suggesting their potential as pectin degraders (García & Chafla, 2015; Hurtado & Otálvaro, 2020; Oumer & Abate, 2018).

A higher number of microorganisms with potential pectinase activity (43.9%) was reported from the Cajamarca region, including the three species *L. xylanilyticus*, *S. maltophilia*, and *S. pavanii*, which might explain the high-quality coffee produced in this region according to the International Specialty Coffee Fair awards obtained in 2018 and 2019 (Agencia Peruana de Noticias, 2019; Cafelab.pe, 2018). These findings provide a starting point for further exploration and quantification of the enzyme activity of indigenous microorganisms from northern Peru and subsequent analyses should be conducted to develop starter cultures of coffee fermentation.

The increasing market demand for coffee consumption has led to the study of indigenous microbiota associated with coffee cultivars to understand the role of these microorganisms in the organoleptic characteristics of the final beverage (Haile & Kang, 2019; Silva et al., 2008a). Indigenous microorganisms with relevant enzyme activity might have the potential to be used as starter cultures, thus allowing controlled fermentation and enhancing the quality of coffee (Carvalho et al., 2017). This scenario might increase the global value chain of coffee and improve the economic incomes of coffee farmers (Melo-Pereira et al., 2017). Therefore, the isolation, identification, and molecular and physicochemical characterization of indigenous microorganisms are essential to accomplish these goals.

## 5 Conclusions

In this study, the microbiota inhabiting wet coffee fermentation from two regions in northern Peru were evaluated. Based on Sanger sequencing and initial screening of pectinolytic activity, we observed that the species composition of fungi and bacteria varied from one coffee farm to another (Amazonas vs. Cajamarca regions). Fungal and bacterial populations were more diverse and abundant in Cajamarca than in the Amazon region. This probably resulted from the longer fermentation time of fruits in Cajamarca that were subjected to a longer degradation and the prevalence of anaerobic microorganisms. The ability of naturally occurring microbiota to degrade mucilage is still unexplored in northern Peru, thereby limiting the development of starter cultures to reduce fermentation times, improve process control, and enhance cup quality. Here, we identify three bacterial species with potential pectinolytic activity (*L. xylanilyticus*, *S. maltophilia* and *S. pavanii*) isolated from the Cajamarca region. This finding should be further investigated by quantifying enzymes activity, performing other biochemical properties of these species (i.e., carbohydrate assimilation, ethanol resistance), and prototyping starter cultures (as a single species or in a consortium) according to coffee variety, local geographic conditions, and fermentation time.

## Funding

This study was funded by the Peruvian National Council for Science and Technology (CONCYTEC) through the Metacafé Project N° 030-2018-FONDECYT-BM-IADT-MU and Proyecto SNIP N° 352439: “Creación de los servicios del centro de investigación, innovación y transferencia tecnológica de café de la Universidad Nacional Toribio Rodríguez de Mendoza, Región de Amazonas” – CEINCAFÉ.

## Acknowledgements

We are most grateful to Mr. Franklin Jhuniór Chinguel Morales from the farm “El Laurel”, Mr. Damian Espinoza Garcia from the farm “La Palma” in the Cajamarca Region, Mr. Denis Huaman from the farm “El Paraíso”, Mr. Job Torres Suarez from the farm “Bombon”, and Mr. Alfonso Tejada Iberico from the farm “Timbuyacu” from the Amazonas region for their valuable help in sample collection. We also thank Stefhany Valdeiglesias Ichillumpa and Ananco Ahuananchi Oswaldo for the translations of the Abstract to Quechua and Awajun languages, respectively.

## References

- Abdollahzadeh, R., Pazhang, M., Najavand, S., Fallahzadeh-Mamaghani, V., & Amani-Ghadim, A. R. (2020). Screening of pectinase-producing bacteria from farmlands and optimization of enzyme production from selected strain by RSM. *Folia Microbiologica*, 65(4), 705-719. <http://dx.doi.org/10.1007/s12223-020-00776-7>. PMID:32026289.
- Agate, A. D., & Bhat, J. V. (1966). Role of pectinolytic yeasts in the degradation of mucilage layer of *Coffea robusta* cherries. *Applied Microbiology*, 14(2), 256-260. <http://dx.doi.org/10.1128/am.14.2.256-260.1966>. PMID:5959859.
- Agencia Peruana de Noticias – ANDINA. (2019, October 26). *Ficafé 2019: meet the winners of Cup of Excellence Peru competition*. Retrieved from <https://www.andina.pe/agencia/noticia-ficafe-2019-conoce-a-los-ganadores-competicion-taza-excelencia-peru-771030.aspx>
- Barragán, J. C. A., Zerpa, S. A. I., Castillo, M. L. S., Haro, I. M. R., Gutiérrez, W. N. A., & Álvarez, F. O. G. (2014). Efecto de la temperatura y pH sobre la actividad y estabilidad de pectinasas producidas por *Bacillus* spp. *REBIOL*, 34(1), 33-41. Retrieved from <https://dialnet.unirioja.es/servlet/articulo?codigo=8143134>
- Beg, Q. K., Bhushan, B., Kapoor, M., & Hoondal, G. S. (2000). Production and characterization of thermostable xylanase and pectinase from *Streptomyces* sp. QG-11-3. *Journal of Industrial Microbiology & Biotechnology*, 24(6), 396-402. <http://dx.doi.org/10.1038/sj.jim.7000010>.
- Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C. H., Xie, D., Suchard, M. A., Rambaut, A., & Drummond, A. J. (2014). BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Computational Biology*, 10(4), e1003537. <http://dx.doi.org/10.1371/journal.pcbi.1003537>. PMID:24722319.
- Bustamante, D. E., Calderon, M. S., Leiva, S., Mendoza, J. E., Arce, M., & Oliva, M. (2021). Three new species of *Trichoderma* in the Harzianum and Longibrachiatum lineages from Peruvian cacao crop soils based on an integrative approach. *Mycologia*, 113(5), 1056-1072. <http://dx.doi.org/10.1080/00275514.2021.1917243>. PMID:34128770.
- Bustamante, D. E., Oliva, M., Leiva, S., Mendoza, J. E., Bobadilla, L., Angulo, G., & Calderon, M. S. (2019). Phylogeny and species delimitations in the entomopathogenic genus *Beauveria* (Hypocreales, Ascomycota), including the description of *B. peruviensis* sp. nov.



- MycKeys*, 58(1), 47-68. <http://dx.doi.org/10.3897/mycokeys.58.35764>. PMID:31565026.
- Cafelab.pe (2018, October 13). *Cup of Excellence Peru 2018: the best coffee is from Cusco*. Retrieved from <https://cafelab.pe/taza-de-excelencia-peru-2018-el-mejor-cafe-es-del-cusco/>
- Carrim, A. J. I., Barbosa, E. C., & Vieira, J. D. G. (2006). Enzymatic activity of endophytic bacterial isolates of *Jacaranda decurrens* Cham. (Carobinha-do-campo). *Brazilian Archives of Biology and Technology*, 49(3), 353-359. <http://dx.doi.org/10.1590/S1516-89132006000400001>.
- Carvalho, D. P. No., de Melo Pereira, G., Tanobe, V., Thomaz Soccol, V., G. da Silva, B., Rodrigues, C., & Soccol, C. (2017). Yeast Diversity and Physicochemical Characteristics Associated with Coffee Bean Fermentation from the Brazilian Cerrado Mineiro Region. *Fermentation (Basel, Switzerland)*, 3(1), 11. <http://dx.doi.org/10.3390/fermentation3010011>.
- Chalfoun, S. M., Cunha, R. L., Carvalho, V. L., & Nogueira, D. A. (2007). Seletividade de fungicidas cúpricos e sistêmicos sobre o fungo *Cladosporium cladosporioides* em cafeeiro. *Summa Phytopathologica*, 33(1), 93-95. <http://dx.doi.org/10.1590/S0100-54052007000100016>.
- Chaudhri, A., & Suneetha, V. (2012). Microbially derived pectinases: a review. *IOSR Journal of Pharmacy and Biological Sciences*, 2(2), 1-5. <http://dx.doi.org/10.9790/3008-0220105>.
- Combo, A. M. M., Aguedo, M., Goffin, D., Wathélet, B., & Paquot, M. (2012). Enzymatic production of pectic oligosaccharides from polygalacturonic acid with commercial pectinase preparations. *Food and Bioprocess Processing*, 90(3), 588-596. <http://dx.doi.org/10.1016/j.fbp.2011.09.003>.
- Cruz-O'Byrne, R., Piraneque-Gambasica, N., & Aguirre-Forero, S. (2021). Microbial diversity associated with spontaneous coffee bean fermentation process and specialty coffee production in northern Colombia. *International Journal of Food Microbiology*, 354(1), 109282. <http://dx.doi.org/10.1016/j.ijfoodmicro.2021.109282>. PMID:34140187.
- De Bruyn, F., Zhang, S. J., Pothakos, V., Torres, J., Lambot, C., Moroni, A. V., Callanan, M., Sybesma, W., Weckx, S., & De Vuyst, L. (2016). Exploring the impact of post-harvest processing on the microbiota and metabolite profiles during a case of green coffee bean production. *Applied and Environmental Microbiology*, 83(1), 1-16. <http://dx.doi.org/10.1128/AEM.02398-16>. PMID:27793826.
- Duong, B., Marraccini, P., Maeght, J. L., Vaast, P., Lebrun, M., & Duponnois, R. (2020). Coffee microbiota and its potential use in sustainable crop management. A review. *Frontiers in Sustainable Food Systems*, 4, 1-31. <http://dx.doi.org/10.3389/fsufs.2020.607935>.
- Elhalis, H., Cox, J., & Zhao, J. (2020). Ecological diversity, evolution and metabolism of microbial communities in the wet fermentation of Australian coffee beans. *International Journal of Food Microbiology*, 321(1), 108544. <http://dx.doi.org/10.1016/j.ijfoodmicro.2020.108544>. PMID:32086129.
- Evangelista, S. R., Miguel, M., Silva, C., Pinheiro, A., & Schwan, R. (2015). Microbiological diversity associated with the spontaneous wet method of coffee fermentation. *International Journal of Food Microbiology*, 210(1), 102-112. <http://dx.doi.org/10.1016/j.ijfoodmicro.2015.06.008>. PMID:26119187.
- Evangelista, S. R., Silva, C. F., da Cruz Miguel, M. G. P., de Souza Cordeiro, C., Pinheiro, A. C. M., Duarte, W. F., & Schwan, R. F. (2014). Improvement of coffee beverage quality by using selected yeasts strains during the fermentation in dry process. *Food Research International*, 61(1), 183-195. <http://dx.doi.org/10.1016/j.foodres.2013.11.033>.
- Feng, X., Dong, H., Yang, P., Yang, R., Lu, J., Lv, J., & Sheng, J. (2016). Culture-dependent and-independent methods to investigate the predominant microorganisms associated with wet processed coffee. *Current Microbiology*, 73(2), 190-195. <http://dx.doi.org/10.1007/s00284-016-1047-3>. PMID:27113591.
- García, C. G., & Chafla, G. A. (2015). *Caracterización de pectinasas antárticas y su uso en la clarificación de jugo de manzana* (Bachelor's thesis). Escuela Superior Politécnica del Litoral, Ecuador. Retrieved from <http://www.dspace.espol.edu.ec/xmlui/handle/123456789/29892>
- García, V., Brizzio, S., Libkind, D., Buzzini, P., & Van Broock, M. (2007). Biodiversity of cold-adapted yeasts from glacial meltwater rivers in Patagonia, Argentina. *FEMS Microbiology Ecology*, 59(2), 331-341. <http://dx.doi.org/10.1111/j.1574-6941.2006.00239.x>. PMID:17313582.
- Garofalo, C., El Khoury, M., Lucas, P., Bely, M., Russo, P., Spano, G., & Capozzi, V. (2015). Autochthonous starter cultures and indigenous grape variety for regional wine production. *Journal of Applied Microbiology*, 118(6), 1395-1408. <http://dx.doi.org/10.1111/jam.12789>. PMID:25752218.
- Haile, M., & Kang, W. H. (2019). The role of microbes in coffee fermentation and their impact on coffee quality. *Journal of Food Quality*, 2019, 4836709. <http://dx.doi.org/10.1155/2019/4836709>.
- Hamdouche, Y., Meile, J. C., Nganou, D. N., Durand, N., Teyssier, C., & Montet, D. (2016). Discrimination of post-harvest coffee processing methods by microbial ecology analyses. *Food Control*, 65(1), 112-120. <http://dx.doi.org/10.1016/j.foodcont.2016.01.022>.
- Hankin, L., & Lacy, G. (1984). Pectinolytic microorganisms. In C. Vanderzant & D. Splittstoesser (Eds.), *Compendium of methods for the microbiological examination of foods* (pp. 176-183). Washington, USA: Speck, ML.
- Huang, C., Becker, M. F., Keto, J. W., & Kovar, D. (2007). Annealing of nanostructured silver films produced by supersonic deposition of nanoparticles. *Journal of Applied Physics*, 102(5), 054308. <http://dx.doi.org/10.1063/1.2776163>.
- Huch, M., & Franz, C. (2015). Coffee. In W. Holzapfel (Ed.), *Advances in fermented foods and beverages: improving quality, technologies and health benefits* (pp. 501-513). Sawston, UK: Woodhead Publishing. <http://dx.doi.org/10.1016/B978-1-78242-015-6.00021-9>.
- Hurtado, G. M., & Otálvaro, H. D. V. (2020). *Determinación de actividad pectinolítica en microbiota levaduriforme aislada a partir de suelos y frutos del Valle del Cauca* (Tesis doctoral). Universidad Santiago de Cali, Cali. Retrieved from <https://repository.usc.edu.co/handle/20.500.12421/3523>
- Joosten, H. M. L. J., Goetz, J., Pittet, A., Schellenberg, M., & Bucheli, P. (2001). Production of ochratoxin A by *Aspergillus carbonarius* on coffee cherries. *International Journal of Food Microbiology*, 65(1-2), 39-44. [http://dx.doi.org/10.1016/S0168-1605\(00\)00506-7](http://dx.doi.org/10.1016/S0168-1605(00)00506-7). PMID:11322699.
- Kashyap, D. R., Chandra, S., Kaul, A., & Tewari, R. (2000). Production, purification and characterization of pectinase from a *Bacillus* sp. DT7. *World Journal of Microbiology & Biotechnology*, 16(3), 277-282. <http://dx.doi.org/10.1023/A:1008902107929>.
- Körner, C. (2007). The use of 'altitude' in ecological research. *Trends in Ecology & Evolution*, 22(11), 569-574. <http://dx.doi.org/10.1016/j.tree.2007.09.006>. PMID:17988759.
- Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33(7), 1870-1874. <http://dx.doi.org/10.1093/molbev/msw054>. PMID:27004904.
- Lanfear, R., Calcott, B., Ho, S. Y., & Guindon, S. (2012). PartitionFinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution*, 29(6), 1695-1701. <http://dx.doi.org/10.1093/molbev/mss020>. PMID:22319168.

- Lechner, S., Mayr, R., Francis, K. P., Prüss, B. M., Kaplan, T., Wiessner-Gunkel, E., Stewart, G. S., & Scherer, S. (1998). *Bacillus weihenstephanensis* sp. nov. is a new psychrotolerant species of the *Bacillus cereus* group. *International Journal of Systematic Bacteriology*, 48(Pt 4), 1373-1382. <http://dx.doi.org/10.1099/00207713-48-4-1373>. PMID:9828439.
- Mantle, P. G., & Chow, A. M. (2000). Ochratoxin formation in *Aspergillus ochraceus* with particular reference to spoilage of coffee. *International Journal of Food Microbiology*, 56(1), 105-109. [http://dx.doi.org/10.1016/S0168-1605\(00\)00278-6](http://dx.doi.org/10.1016/S0168-1605(00)00278-6). PMID:10857930.
- Martins, P. M. M., Batista, N. N., Miguel, M. G. D. C. P., Simão, J. B. P., Soares, J. R., & Schwan, R. F. (2020). Coffee growing altitude influences the microbiota, chemical compounds and the quality of fermented coffees. *Food Research International*, 129(1), 108872. <http://dx.doi.org/10.1016/j.foodres.2019.108872>. PMID:32036899.
- Masoud, W., & Kaltoft, C. H. (2006). The effects of yeasts involved in the fermentation of *Coffea arabica* in East Africa on growth and ochratoxin A (OTA) production by *Aspergillus ochraceus*. *International Journal of Food Microbiology*, 106(2), 229-234. <http://dx.doi.org/10.1016/j.ijfoodmicro.2005.06.015>. PMID:16213049.
- Masoud, W., Cesar, L. B., Jespersen, L., & Jakobsen, M. (2004). Yeast involved in fermentation of *Coffea arabica* in East Africa determined by genotyping and by direct denaturing gradient gel electrophoresis. *Yeast (Chichester, England)*, 21(7), 549-556. <http://dx.doi.org/10.1002/yea.1124>. PMID:15164358.
- Melo Pereira, G. V., Soccol, V. T., Pandey, A., Medeiros, A. B., Andrade Lara, J. M., Gollo, A. L., & Soccol, C. R. (2014). Isolation, selection and evaluation of yeasts for use in fermentation of coffee beans by the wet process. *International Journal of Food Microbiology*, 188(1), 60-66. <http://dx.doi.org/10.1016/j.ijfoodmicro.2014.07.008>. PMID:25087206.
- Melo-Pereira, G. V., Soccol, V. T., Brar, S. K., Neto, E., & Soccol, C. R. (2017). Microbial ecology and starter culture technology in coffee processing. *Critical Reviews in Food Science and Nutrition*, 57(13), 2775-2788. <http://dx.doi.org/10.1080/10408398.2015.1067759>. PMID:26462969.
- Murad, H. A., & Azzaz, H. H. (2011). Microbial pectinases and ruminant nutrition. *Research Journal of Microbiology*, 6(3), 246-269. <http://dx.doi.org/10.3923/jm.2011.246.269>.
- Oktaviani, L., Astuti, D. I., Rosmiati, M., & Abduh, M. Y. (2020). Fermentation of coffee pulp using indigenous lactic acid bacteria with simultaneous aeration to produce cascara with a high antioxidant activity. *Heliyon*, 6(7), e04462. <http://dx.doi.org/10.1016/j.heliyon.2020.e04462>. PMID:32743093.
- Oliveira Junqueira, A. C., de Melo Pereira, G. V., Coral Medina, J. D., Alvear, M. C. R., Rosero, R., de Carvalho D. P. No., Enríquez, H. G., & Soccol, C. R. (2019). First description of bacterial and fungal communities in Colombian coffee beans fermentation analysed using Illumina-based amplicon sequencing. *Scientific Reports*, 9(1), 8794. <http://dx.doi.org/10.1038/s41598-019-45002-8>. PMID:31217528.
- Oliveira, M. N., Santos, T. M., Vale, H. M., Delvaux, J. C., Cordero, A. P., Ferreira, A. B., Miguel, P. S., Tótola, M. R., Costa, M. D., Moraes, C. A., & Borges, A. C. (2013). Endophytic microbial diversity in coffee cherries of *Coffea arabica* from southeastern Brazil. *Canadian Journal of Microbiology*, 59(4), 221-230. <http://dx.doi.org/10.1139/cjm-2012-0674>. PMID:23586745.
- Oumer, O. J., & Abate, D. (2018). Screening and molecular identification of pectinase producing microbes from coffee pulp. *BioMed Research International*, 2018, 2961767. <http://dx.doi.org/10.1155/2018/2961767>. PMID:29850500.
- Patidar, M. K., Nighojkar, S., Kumar, A., & Nighojkar, A. (2018). Pectinolytic enzymes-solid state fermentation, assay methods and applications in fruit juice industries: a review. *3 Biotech*, 8(4), 199. <https://doi.org/10.1007/s13205-018-1220-4>.
- Pereira, R. T. G., Pfenning, L. H., & Castro, H. A. (2005). Caracterização e dinâmica de colonização de *Cladosporium cladosporioides* (Fresen.) de vries em frutos do cafeeiro (*Coffea arabica* L.). *Ciência e Agrotecnologia*, 29(6), 1112-1116. <http://dx.doi.org/10.1590/S1413-70542005000600002>.
- Pothakos, V., De Vuyst, L., Zhang, S. J., De Bruyn, F., Verce, M., Torres, J., Callanan, M., Moccand, C., & Weckx, S. (2020). Temporal shotgun metagenomics of an Ecuadorian coffee fermentation process highlights the predominance of lactic acid bacteria. *Current Research in Biotechnology*, 2(1), 1-15. <http://dx.doi.org/10.1016/j.crbiot.2020.02.001>.
- Poveda, G., Gil-Durán, C., Vaca, I., Levicán, G., & Chávez, R. (2018). Cold-active pectinolytic activity produced by filamentous fungi associated with Antarctic marine sponges. *Biological Research*, 51(1), 28. <http://dx.doi.org/10.1186/s40659-018-0177-4>. PMID:30149803.
- Puerta, G. I., & Echeverry, J. G. (2015). *Fermentación controlada del café: Tecnología para agregar valor a la calidad* (Avances Técnicos, No. 454, 12 p.). Manizales: CENICAFE. Retrieved from <http://hdl.handle.net/10778/558>
- Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D. L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M. A., & Huelsenbeck, J. P. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology*, 61(3), 539-542. <http://dx.doi.org/10.1093/sysbio/sys029>. PMID:22357727.
- Roussos, S., de los Angeles Aquíhuatl, M., del Refugio Trejo-Hernández, M., Gaime Perraud, I., Favela, E., Ramakrishna, M., Raimbault, M., & Viniegra-González, G. (1995). Biotechnological management of coffee pulp-isolation, screening, characterization, selection of caffeine-degrading fungi and natural microflora present in coffee pulp and husk. *Applied Microbiology and Biotechnology*, 42(5), 756-762. <http://dx.doi.org/10.1007/BF00171958>.
- Santos, A., van Aerle, R., Barrientos, L., & Martínez-Urtaza, J. (2020). Computational methods for 16S metabarcoding studies using Nanopore sequencing data. *Computational and Structural Biotechnology Journal*, 18(1), 296-305. <http://dx.doi.org/10.1016/j.csbj.2020.01.005>. PMID:32071706.
- Schwan, R. S., Silva, V., Ferreira, C., & Batista, L. R. (2012). Coffee fermentation. In E. O. Evranuz (Ed.), *Handbook of plant-based fermented food and beverage technology* (pp. 677-690). Boca Raton, FL: CRC Press. <http://dx.doi.org/10.1201/b12055-49>.
- Schwan, R., Cooper, R., & Wheals, A. (1997). Endopolygalacturonase secretion by *Kluyveromyces marxianus* and other cocoa pulp-degrading yeasts. *Enzyme and Microbial Technology*, 21(4), 234-244. [http://dx.doi.org/10.1016/S0141-0229\(96\)00261-X](http://dx.doi.org/10.1016/S0141-0229(96)00261-X).
- Silva, B. L., Pereira, P. V., Bertoli, L. D., Silveira, D. L., Batista, N. N., Pinheiro, P. F., de Souza Carneiro, J., Schwan, R. F., de Assis Silva, S., Coelho, J. M., & Bernardes, P. C. (2021). Fermentation of *Coffea canephora* inoculated with yeasts: Microbiological, chemical, and sensory characteristics. *Food Microbiology*, 98(1), 103786. <http://dx.doi.org/10.1016/j.fm.2021.103786>. PMID:33875214.
- Silva, C. F., Schwan, R. F., Dias, E. S., & Wheals, A. E. (2000). Microbial diversity during maturation and natural processing of coffee cherries of *Coffea arabica* in Brazil. *International Journal of Food Microbiology*, 60(2-3), 251-260. [http://dx.doi.org/10.1016/S0168-1605\(00\)00315-9](http://dx.doi.org/10.1016/S0168-1605(00)00315-9). PMID:11016614.
- Silva, C. F., Batista, L. R., Abreu, L. M., Dias, E. S., & Schwan, R. F. (2008a). Succession of bacterial and fungal communities during natural coffee (*Coffea arabica*) fermentation. *Food Microbiology*, 25(8), 951-957. <http://dx.doi.org/10.1016/j.fm.2008.07.003>. PMID:18954729.

- Silva, C. F., Batista, L. R., & Schwan, R. F. (2008b). Incidence and distribution of filamentous fungi during fermentation, drying and storage of coffee (*Coffea arabica* L.) beans. *Brazilian Journal of Microbiology*, 39(3), 521-526. <http://dx.doi.org/10.1590/S1517-83822008000300022>. PMID:24031259.
- Silva, C. F., Vilela, D. M., de Souza Cordeiro, C., Duarte, W. F., Dias, D. R., & Schwan, R. F. (2013). Evaluation of a potential starter culture for enhance quality of coffee fermentation. *World Journal of Microbiology & Biotechnology*, 29(2), 235-247. <http://dx.doi.org/10.1007/s11274-012-1175-2>. PMID:23054699.
- Silva, E. G., Borges, M. D. F., Medina, C., Piccoli, R. H., & Schwan, R. F. (2005). Pectinolytic enzymes secreted by yeasts from tropical fruits. *FEMS Yeast Research*, 5(9), 859-865. <http://dx.doi.org/10.1016/j.femsyr.2005.02.006>. PMID:15925314.
- Silvestro, D., & Michalak, I. (2012). raxmlGUI: a graphical front-end for RAxML. *Organisms, Diversity & Evolution*, 12(4), 335-337. <http://dx.doi.org/10.1007/s13127-011-0056-0>.
- Souza, J. V., Silva, É. S., Maia, M. L., & Teixeira, M. F. (2003). Screening of fungal strains for pectinolytic activity: endopolygalacturonase production by *Peacilomyces clavissporus* 2A. UMIDA. 1. *Process Biochemistry*, 39(4), 455-458. [http://dx.doi.org/10.1016/S0032-9592\(03\)00092-X](http://dx.doi.org/10.1016/S0032-9592(03)00092-X).
- Souza, M. L. D., Passamani, F. R. F., Ávila, C. L. D. S., Batista, L. R., Schwan, R. F., & Silva, C. F. (2017). Use of wild yeasts as a biocontrol agent against toxigenic fungi and OTA production. *Acta Scientiarum. Agronomy*, 39(1), 349-358. <http://dx.doi.org/10.4025/actasciagron.v39i3.32659>.
- Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics (Oxford, England)*, 30(9), 1312-1313. <http://dx.doi.org/10.1093/bioinformatics/btu033>. PMID:24451623.
- Sunnotel, O., & Nigam, P. (2002). Pectinolytic activity of bacteria isolated from soil and two fungal strains during submerged fermentation. *World Journal of Microbiology & Biotechnology*, 18(9), 835-839. <http://dx.doi.org/10.1023/A:1021209123641>.
- Technelysium. (1998). *Chromas*. Retrieved from <http://www.technelysium.com.au/chromas.html>
- Teshome, B., Wassie, M., & Abatneh, E. (2017). Isolation, screening and biochemical characterization of phosphate solubilizing rhizobacteria associated with *Coffea arabica* L. *Journal of Fertilizers & Pesticides*, 8(03). <http://dx.doi.org/10.4172/2471-2728.1000188>.
- Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22(22), 4673-4680. <http://dx.doi.org/10.1093/nar/22.22.4673>. PMID:7984417.
- Vera, K., Valdivieso, W., Mantilla, A., Jaimes, W., Torrado, J., & Zafra, G. (2018). Functional Metagenomic Analysis of the Coffee (*Coffea arabica*) Fermentation. *Chemical Engineering Transactions*, 64(1), 355-360. <http://dx.doi.org/10.3303/CET1864060>.
- Vilela, D. M., Pereira, G. V. D. M., Silva, C. F., Batista, L. R., & Schwan, R. F. (2010). Molecular ecology and polyphasic characterization of the microbiota associated with semi-dry processed coffee (*Coffea arabica* L.). *Food Microbiology*, 27(8), 1128-1135. <http://dx.doi.org/10.1016/j.fm.2010.07.024>. PMID:20832694.
- Wang, L., & Stegemann, J. P. (2010). Extraction of high quality RNA from polysaccharide matrices using cetlytrimethylammonium bromide. *Biomaterials*, 31(7), 1612-1618. <http://dx.doi.org/10.1016/j.biomaterials.2009.11.024>. PMID:19962190.
- Wood, W., & Kellogg, S. (1988). *Methods in enzymology* (Vol. 161: Biomass, Part B: Lignin, Pectin, and Chitin, pp. 315-322). New York: Academic Press.
- Xavier-Santos, S., Carvalho, C., Bonfá, M., Silva, R., Capelari, M., & Gomes, E. (2004). Screening for pectinolytic activity of wood-rotting basidiomycetes and characterization of the enzymes. *Folia Microbiologica*, 49(1), 46-52. <http://dx.doi.org/10.1007/BF02931645>. PMID:15114865.
- Zhang, S. J., De Bruyn, F., Pothakos, V., Torres, J., Falconi, C., Moccand, C., Weckx, S., & de Vuyst, L. (2019). Following coffee production from cherries to cup: microbiological and metabolomic analysis of wet processing of *Coffea arabica*. *Applied and Environmental Microbiology*, 85(6), e02635-e18. <http://dx.doi.org/10.1128/AEM.02635-18>. PMID:30709820.
- Zhao, L., Xu, Y., Lai, X. H., Shan, C., Deng, Z., & Ji, Y. (2015). Screening and characterization of endophytic *Bacillus* and *Paenibacillus* strains from medicinal plant *Lonicera japonica* for use as potential plant growth promoters. *Brazilian Journal of Microbiology*, 46(4), 977-989. <http://dx.doi.org/10.1590/S1517-838246420140024>. PMID:26691455.



## Supplementary material

Supplementary material accompanies this paper.

**Table S1.** List of taxa used in the molecular analyses along with strain and, if known, country. (T) represents sequence obtained from type strains. GenBank accession numbers under each marker; if marker not sequenced indicated by “-”. Sequences generated in present study are in bold.

**Table S2.** Best models for maximum likelihood and Bayesian inferences by group and taxonomical class.

**Figure S1.** Number of species per taxonomical class for bacteria and fungi isolated from coffee ferments in Amazonas and Cajamarca Regions.

**Figure S2.** Phylogenetic tree of the Actinobacteria class based on maximum likelihood inferences of 16S rRNA gene sequences. Bootstrap values of maximum likelihood (BS;  $\geq 50\%$ )/Bayesian posterior probabilities (BPP;  $\geq 0.9$ ) are indicated above the branches. Values below 50% (BS) or 0.90 (BPP) are indicated with dashes (-). The scale bar indicates the number of nucleotide substitutions per site. Sequence accession numbers are shown in parentheses, with T representing sequences obtained from type strains. Sequences for taxa in bold were generated in this study.

**Figure S3.** Phylogenetic tree of class Alphaproteobacteria based on maximum likelihood inference of 16S rRNA gene sequences. Bootstrap values of maximum likelihood (BS;  $\geq 50\%$ )/Bayesian posterior probabilities (BPP;  $\geq 0.9$ ) are indicated above the branches. Values below 50% (BS) or 0.90 (BPP) are indicated with dashes (-). The scale bar indicates the number of nucleotide substitutions per site. Sequence accession numbers are shown in parentheses, with T representing sequences obtained from type strains. Sequences for taxa in bold were generated in this study.

**Figure S4.** Phylogenetic tree of class Betaproteobacteria based on maximum likelihood inference of 16S rRNA gene sequences. Bootstrap values of maximum likelihood (BS;  $\geq 50\%$ )/Bayesian posterior probabilities (BPP;  $\geq 0.9$ ) are indicated above the branches. Values below 50% (BS) or 0.90 (BPP) are indicated with dashes (-). The scale bar indicates the number of nucleotide substitutions per site. Sequence accession numbers are shown in parentheses, with T representing sequences obtained from type strains. Sequences for taxa in bold were generated in this study.

**Figure S5.** Phylogenetic tree of the class Firmicutes based on maximum likelihood inference of 16S rRNA gene sequences. Bootstrap values of maximum likelihood (BS;  $\geq 50\%$ )/Bayesian posterior probabilities (BPP;  $\geq 0.9$ ) are indicated above the branches. Values below 50% (BS) or 0.90 (BPP) are indicated with dashes (-). The scale bar indicates the number of nucleotide substitutions per site. Sequence accession numbers are shown in parentheses, with T representing sequences obtained from type strains. Sequences for taxa in bold were generated in this study.

**Figure S6.** Phylogenetic tree of class Gammaproteobacteria based on maximum likelihood inference of 16S rRNA gene sequences. Bootstrap values of maximum likelihood (BS;  $\geq 50\%$ )/Bayesian posterior probabilities (BPP;  $\geq 0.9$ ) are indicated above the branches. Values below 50% (BS) or 0.90 (BPP) are indicated with dashes (-). The scale bar indicates the number of nucleotide substitutions per site. Sequence accession numbers are shown in parentheses, with T representing sequences obtained from type strains. Sequences for taxa in bold were generated in this study.

**Figure S7.** Phylogenetic tree of class Dothideomycetes based on maximum likelihood inference of ITS sequences. Bootstrap values of maximum likelihood (BS;  $\geq 50\%$ )/Bayesian posterior probabilities (BPP;  $\geq 0.9$ ) are indicated above the branches. Values below 50% (BS) or 0.90 (BPP) are indicated with dashes (-). The scale bar indicates the number of nucleotide substitutions per site. Sequence accession numbers are shown in parentheses, with T representing sequences obtained from type strains. Sequences for taxa in bold were generated in this study.

**Figure S8.** Phylogenetic tree of class Eurotiomycetes based on maximum likelihood inference of ITS sequences. Bootstrap values of maximum likelihood (BS;  $\geq 50\%$ )/Bayesian posterior probabilities (BPP;  $\geq 0.9$ ) are indicated above the branches. Values below 50% (BS) or 0.90 (BPP) are indicated with dashes (-). The scale bar indicates the number of nucleotide substitutions per site. Sequence accession numbers are shown in parentheses, with T representing sequences obtained from type strains. Sequences for taxa in bold were generated in this study.

**Figure S9.** Phylogenetic tree of class Sordariomycetes based on maximum likelihood inference of ITS sequences. Bootstrap values of maximum likelihood (BS;  $\geq 50\%$ )/Bayesian posterior probabilities (BPP;  $\geq 0.9$ ) are indicated above the branches. Values below 50% (BS) or 0.90 (BPP) are indicated with dashes (-). The scale bar indicates the number of nucleotide substitutions per site. Sequence accession numbers are shown in parentheses, with T representing sequences obtained from type strains. Sequences for taxa in bold were generated in this study.

**Figure S10.** Microorganisms before and after 1% CTAB precipitation in MP5 and/or MP7 media and the formation of different types of halos. Microorganisms from the Cajamarca Region: *Penicillium citrinum* without CETAB (A, MP5), *Penicillium citrinum* with CETAB (A', MP5), *Aspergillus* sp.1 without CETAB (B, MP7; C, MP7), *Aspergillus* sp.1 with CETAB (B', MP7; C', MP7), *Alcaligenes* sp. without CETAB (D, MP7), *Alcaligenes* sp. with CETAB (D', MP7), *Clostridium algidixylanolyticum* without CETAB (E, MP5), *Clostridium algidixylanolyticum* with CETAB (E', MP5). Microorganisms from the Amazonas region: *Lysinibacillus fusiformis* without CETAB (F, MP7; G, MP7; H, MP5) and *Lysinibacillus* sp. 4 with CETAB (F', MP7; G', MP7; H', MP5).

This material is available as part of the online article from <https://doi.org/10.1590/fst.81922>